

APCTP SEMINAR

Muon ($g-2$) and Thermal WIMP DM in $U(1) L Q -L \tau$ Models

Jongkuk Kim

KIAS

September 14th (Wed.) 16:00

#512, APCTP & Online via ZOOM

$U(1)LQ-L\tau \equiv U(1)X$ model is anomaly free within the Standard Model (SM) fermion content, and can accommodate the muon ($g-2$) data for $MZ' \sim O(10-100)$ MeV and $gX \sim (4-8) \times 10^{-4}$. WIMP type thermal dark matter (DM) can be also introduced for $MZ' \sim 2MDM$, if DM pair annihilations into the SM particles occur only through the s-channel Z' exchange. In this work, we show that this tight correlation between MZ' and MDM can be completely evaded both for scalar and fermionic DM, if we include the contributions from dark Higgs boson. Dark Higgs boson plays a crucial role in DM phenomenology, not only for generation of dark photon mass, but also opening new channels for DM pair annihilations into the final states involving dark Higgs boson, such as dark Higgs pair as well as $Z'Z'$ through dark Higgs exchange in the s-channel, and co-annihilation into $Z'H_1$ in case of inelastic DM. Thus dark Higgs boson will dissect the strong correlation $MZ' \sim 2MDM$, and much wider mass range is allowed for $U(1)X$ -charged complex scalar and Dirac fermion DM, still explaining the muon ($g-2$).

■ ZOOM Webinar

1) Please register through this ZOOM link (password is 0)

<https://us06web.zoom.us/meeting/register/tZlocO6vpjMuEtY7vymbXo07KyfqL4Oashle>

2) Join the webinar with a link generated after the registration

3) Please rename your profile - E.g. **Full name (affiliation)**

■ Contact information

- Host: Hiroshi Okada (hiroshi.okada@apctp.org)

- Office: Research Support Team (ra@apctp.org)